

CBSE Class 12 Maths Question Paper Solution 2018

QUESTION PAPER CODE 65/1

EXPECTED ANSWER/VALUE POINTS

SECTION A $\frac{1}{2} + \frac{1}{2}$

1. $\frac{\pi}{3} - \left(\pi - \frac{\pi}{6} \right) = -\frac{\pi}{2}$

Note: $\frac{1}{2}$ m. for any one of the two correct values and $\frac{1}{2}$ m. for final answer

2. $a = -2, b = 3$ $\frac{1}{2} + \frac{1}{2}$

3. $|\vec{a}| = |\vec{b}| = 3$ $\frac{1}{2} + \frac{1}{2}$

4. $5 \circ 10 = (5 * 10) + 3 = 10 + 3 = 13$ For $5 * 10 = 10$ $\frac{1}{2}$

For Final Answer = 13 $\frac{1}{2}$

SECTION B

5. In RHS, put $x = \sin \theta$ $\frac{1}{2}$

$$\begin{aligned} \text{RHS} &= \sin^{-1} (3 \sin \theta - 4 \sin^3 \theta) \\ &= \sin^{-1} (\sin 3\theta) \end{aligned} \quad \text{1}$$

$$= 3\theta = 3 \sin^{-1} x = \text{LHS.} \quad \text{1}$$

6. $|A| = 2, \therefore A^{-1} = \frac{1}{2} \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$ 1

$$\text{LHS} = 2A^{-1} = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}, \text{ RHS} = 9 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix} = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix} \quad \text{1}$$

$\therefore \text{LHS} = \text{RHS}$

7. $f(x) = \tan^{-1}\left(\frac{1+\cos x}{\sin x}\right) = \tan^{-1}\left(\frac{2\cos^2 x/2}{2\sin x/2 \cos x/2}\right)$ 1

$$= \tan^{-1}\left(\cot x/2\right) = \frac{\pi}{2} - \frac{x}{2}$$
 $\frac{1}{2}$

$$\therefore f'(x) = -\frac{1}{2}$$
 $\frac{1}{2}$

8. Marginal cost = $C'(x) = 0.015x^2 - 0.04x + 30$ 1

At $x = 3$, $C'(3) = 30.015$ 1

9. $I = \int \frac{1-2\sin^2 x+2\sin^2 x}{\cos^2 x} dx$ $\frac{1}{2}$

$$= \int \sec^2 x dx$$
 1

$$= \tan x + C$$
 $\frac{1}{2}$

10. $\frac{dy}{dx} = bae^{bx+5} \Rightarrow \frac{dy}{dx} = by$ 1

$$\Rightarrow \frac{d^2y}{dx^2} = b \frac{dy}{dx}$$
 $\frac{1}{2}$

\therefore The differential equation is: $y \frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$ $\frac{1}{2}$

11. $\sin \theta = \frac{|(\hat{i} - 2\hat{j} + 3\hat{k}) \times (3\hat{i} - 2\hat{j} + \hat{k})|}{|\hat{i} - 2\hat{j} + 3\hat{k}| |3\hat{i} - 2\hat{j} + \hat{k}|}$ $\frac{1}{2}$

$$|(\hat{i} - 2\hat{j} + 3\hat{k}) \times (3\hat{i} - 2\hat{j} + \hat{k})| = |4\hat{i} + 8\hat{j} + 4\hat{k}| = 4\sqrt{6}$$
 1

$$\sin \theta = \frac{4\sqrt{6}}{14} = \frac{2\sqrt{6}}{7}$$
 $\frac{1}{2}$

12. A: Getting a sum of 8, B: Red die resulted in no. < 4

$$P(A/B) = \frac{P(A \cap B)}{P(B)}$$
 1

$$= \frac{2/36}{18/36} = \frac{1}{9}$$
 1

SECTION C

13. LHS =
$$\begin{vmatrix} 1 & 1 & 1+3x \\ 1+3y & 1 & 1 \\ 1 & 1+3z & 1 \end{vmatrix}$$

$$= \begin{vmatrix} 1 & 0 & 3x \\ 1+3y & -3y & -3y \\ 1 & 3z & 0 \end{vmatrix} \quad (\text{Using } C_2 \rightarrow C_2 - C_1 \text{ & } C_3 \rightarrow C_3 - C_1)$$

1+1
(Any two relevant operations)

$$= 1 \times (9yz) + 3x(3z + 9yz + 3y) \quad (\text{Expanding along } R_1)$$

$$= 9(3xyz + xy + yz + zx) = \text{RHS}$$

14. Differentiating with respect to 'x'

$$2(x^2 + y^2) \left(2x + 2y \frac{dy}{dx} \right) = x \frac{dy}{dx} + y$$

2

$$\Rightarrow \frac{dy}{dx} = \frac{y - 4x^3 - 4xy^2}{4x^2y + 4y^3 - x}$$

2

OR

$$\frac{dx}{d\theta} = a(2 - 2 \cos 2\theta) = 4a \sin^2 \theta$$

1

$$\frac{dy}{d\theta} = 2a \sin 2\theta = 4a \sin \theta \cdot \cos \theta$$

1

$$\therefore \frac{dy}{dx} = \frac{4a \sin \theta \cos \theta}{4a \sin^2 \theta} = \cot \theta$$

1

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{3}} = \frac{1}{\sqrt{3}}$$

1

15. $y = \sin(\sin x) \Rightarrow \frac{dy}{dx} = \cos(\sin x) \cdot \cos x$

1

and $\frac{d^2y}{dx^2} = -\sin(\sin x) \cdot \cos^2 x - \sin x \cos(\sin x)$

1+1

$$\text{LHS} = -\sin(\sin x) \cos^2 x - \sin x \cos(\sin x) + \frac{\sin x}{\cos x} \cos(\sin x) \cos x + \sin(\sin x) \cos^2 x$$

1

$$= 0 = \text{RHS}$$

(3)

16. $x_1 = 2 \Rightarrow y_1 = 3 \quad (\because y_1 > 0)$

 $\frac{1}{2}$

Differentiating the given equation, we get, $\frac{dy}{dx} = \frac{-16x}{9y}$

 $\frac{1}{2}$

Slope of tangent at $(2, 3) = \left. \frac{dy}{dx} \right|_{(2, 3)} = -\frac{32}{27}$

 $\frac{1}{2}$

Slope of Normal at $(2, 3) = \frac{27}{32}$

 $\frac{1}{2}$

Equation of tangent: $32x + 27y = 145$

1

Equation of Normal: $27x - 32y = -42$

1

OR

$f'(x) = x^3 - 3x^2 - 10x + 24$

 $\frac{1}{2}$

$= (x-2)(x-4)(x+3)$

1

$f'(x) = 0 \Rightarrow x = -3, 2, 4.$

 $\frac{1}{2}$ sign of $f'(x)$:

$\therefore f(x)$ is strictly increasing on $(-3, 2) \cup (4, \infty)$

1

and $f(x)$ is strictly decreasing on $(-\infty, -3) \cup (2, 4)$

1

17. Let side of base = x and depth of tank = y

$V = x^2y \Rightarrow y = \frac{V}{x^2}, (V = \text{Quantity of water} = \text{constant})$

Cost of material is least when area of sheet used is minimum.

$A(\text{Surface area of tank}) = x^2 + 4xy = x^2 + \frac{4V}{x}$

 $\frac{1}{2} + \frac{1}{2}$

$\frac{dA}{dx} = 2x - \frac{4V}{x^2}, \frac{dA}{dx} = 0 \Rightarrow x^3 = 2V, y = \frac{x^3}{2x^2} = \frac{x}{2}$

 $\frac{1}{2} + \frac{1}{2}$

$\frac{d^2A}{dx^2} = 2 + \frac{8V}{x^3} > 0, \therefore \text{Area is minimum, thus cost is minimum when } y = \frac{x}{2}$

 $\frac{1}{2} + \frac{1}{2}$

Value: Any relevant value.

1

18. Put $\sin x = t \Rightarrow \cos x dx = dt$

$\frac{1}{2}$

$$\text{Let } I = \int \frac{2 \cos x}{(1-\sin x)(1+\sin^2 x)} dx = \int \frac{2}{(1-t)(1+t^2)} dt$$

$$\text{Let } \frac{2}{(1-t)(1+t^2)} = \frac{A}{1-t} + \frac{Bt+C}{1+t^2}, \text{ solving we get}$$

$$A = 1, B = 1, C = 1$$

$1\frac{1}{2}$

$$\therefore I = \int \frac{1}{1-t} dt + \frac{1}{2} \int \frac{2t}{1+t^2} + \int \frac{1}{1+t^2} dt$$

$$= -\log|1-t| + \frac{1}{2} \log|1+t^2| + \tan^{-1} t + C$$

$1\frac{1}{2}$

$$= -\log(1-\sin x) + \frac{1}{2} \log(1+\sin^2 x) + \tan^{-1}(\sin x) + C$$

$\frac{1}{2}$

19. Separating the variables, we get:

$$\int \frac{\sec^2 y}{\tan y} dy = \int \frac{e^x}{e^x - 2} dx$$

$1\frac{1}{2}$

$$\Rightarrow \log|\tan y| = \log|e^x - 2| + \log C$$

$1\frac{1}{2}$

$$\Rightarrow \tan y = C(e^x - 2), \text{ for } x = 0, y = \pi/4, C = -1$$

$\frac{1}{2}$

$$\therefore \text{Particular solution is: } \tan y = 2 - e^x.$$

$\frac{1}{2}$

OR

$$\text{Integrating factor} = e^{\int 2 \tan x dx} = \sec^2 x$$

1

$$\therefore \text{Solution is: } y \cdot \sec^2 x = \int \sin x \cdot \sec^2 x dx = \int \sec x \cdot \tan x dx$$

1

$$\Rightarrow y \cdot \sec^2 x = \sec x + C, \text{ for } x = \frac{\pi}{3}, y = 0, \therefore C = -2$$

$1+\frac{1}{2}$

$$\therefore \text{Particular solution is: } y \cdot \sec^2 x = \sec x - 2$$

$\frac{1}{2}$

$$\text{or } y = \cos x - 2 \cos^2 x$$

20. $\vec{d} = \lambda(\vec{c} \times \vec{b}) = \lambda \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & -1 \\ 1 & -4 & 5 \end{vmatrix}$ 1

$\therefore \vec{d} = \lambda \hat{i} - 16\lambda \hat{j} - 13\lambda \hat{k}$ 1

$\vec{d} \cdot \vec{a} = 21 \Rightarrow 4\lambda - 80\lambda + 13\lambda = 21 \Rightarrow \lambda = -\frac{1}{3}$ 1

$\therefore \vec{d} = -\frac{1}{3}\hat{i} + \frac{16}{3}\hat{j} + \frac{13}{3}\hat{k}$ 1

21. Here $\vec{a}_1 = 4\hat{i} - \hat{j}$, $\vec{a}_2 = \hat{i} - \hat{j} + 2\hat{k}$, $\vec{a}_2 - \vec{a}_1 = -3\hat{i} + 2\hat{k}$ 1

$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -3 \\ 2 & 4 & -5 \end{vmatrix} = 2\hat{i} - \hat{j}$ 1

Shortest distance = $\frac{|(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|}$ 1

$= \left| \frac{-6}{\sqrt{5}} \right| = \frac{6}{\sqrt{5}}$ or $\frac{6\sqrt{5}}{5}$ 1

22. E_1 : She gets 1 or 2 on die.
 E_2 : She gets 3, 4, 5 or 6 on die.
A: She obtained exactly 1 tail

$P(E_1) = \frac{1}{3}$, $P(E_2) = \frac{2}{3}$
 $P(A/E_1) = \frac{3}{8}$, $P(A/E_2) = \frac{1}{2}$

$P(E_2/A) = \frac{P(E_2) \cdot P(A/E_2)}{P(E_1) \cdot P(A/E_1) + P(E_2) \cdot P(A/E_2)}$ 1

$= \frac{\frac{2}{3} \times \frac{1}{2}}{\frac{1}{3} \times \frac{3}{8} + \frac{2}{3} \times \frac{1}{2}} = \frac{8}{11}$ 1

23. Let X denote the larger of two numbers

X	2	3	4	5	$\frac{1}{2}$
$P(X)$	$1/10$	$2/10$	$3/10$	$4/10$	1
$X \cdot P(X)$	$2/10$	$6/10$	$12/10$	$20/10$	$\frac{1}{2}$
$X^2 \cdot P(X)$	$4/10$	$18/10$	$48/10$	$100/10$	$\frac{1}{2}$

$$\text{Mean} = \sum X \cdot P(X) = \frac{40}{10} = 4 \quad \frac{1}{2}$$

$$\text{Variance} = \sum X^2 \cdot P(X) - [\sum X \cdot P(X)]^2 = \frac{170}{10} - 4^2 = 1 \quad 1$$

SECTION D

24. Reflexive: $|a - a| = 0$, which is divisible by 4, $\forall a \in A$

1

$$\therefore (a, a) \in R, \forall a \in A \quad \therefore R \text{ is reflexive}$$

Symmetric: let $(a, b) \in R$

$\Rightarrow |a - b|$ is divisible by 4

$\Rightarrow |b - a|$ is divisible by 4 ($\because |a - b| = |b - a|$)

$\Rightarrow (b, a) \in R \quad \therefore R \text{ is symmetric.}$

1

Transitive: let $(a, b), (b, c) \in R$

$\Rightarrow |a - b| \& |b - c|$ are divisible by 4

$\Rightarrow a - b = \pm 4m, b - c = \pm 4n, m, n \in \mathbb{Z}$

Adding we get, $a - c = 4(\pm m \pm n)$

$\Rightarrow (a - c)$ is divisible by 4

$\Rightarrow |a - c|$ is divisible by 4 $\therefore (a, c) \in R$

$\therefore R$ is transitive

2

Hence R is an equivalence relation in A

1

set of elements related to 1 is $\{1, 5, 9\}$

and $[2] = \{2, 6, 10\}$.

}

1

OR

Here $f(2) = f\left(\frac{1}{2}\right) = \frac{2}{5}$ but $2 \neq \frac{1}{2}$

$\therefore f$ is not 1-1

2

for $y = \frac{1}{\sqrt{2}}$ let $f(x) = \frac{1}{\sqrt{2}} \Rightarrow x^2 - \sqrt{2}x + 1 = 0$

As $D = (-\sqrt{2})^2 - 4(1)(1) < 0$, \therefore No real solution

$\therefore f(x) \neq \frac{1}{\sqrt{2}}$, for any $x \in R(D_f)$ $\therefore f$ is not onto

2

$$\text{fog}(x) = f(2x - 1) = \frac{2x - 1}{(2x - 1)^2 + 1} = \frac{2x - 1}{4x^2 - 4x + 2}$$

2

25. $|A| = -1 \neq 0 \quad \therefore A^{-1}$ exists

1

Co-factors of A are:

$$\begin{aligned} A_{11} &= 0 ; & A_{12} &= 2 ; & A_{13} &= 1 \\ A_{21} &= -1 ; & A_{22} &= -9 ; & A_{23} &= -5 \\ A_{31} &= 2 ; & A_{32} &= 23 ; & A_{33} &= 13 \end{aligned} \quad \left. \begin{array}{l} 1 \text{ m for any} \\ 4 \text{ correct} \\ \text{cofactors} \end{array} \right\}$$

2

$$\text{adj}(A) = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix} \Rightarrow A^{-1} = \frac{1}{|A|} \cdot \text{adj}(A) = \frac{1}{-1} \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix}$$

1

2

For : $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix}$, the system of equation is $A \cdot X = B$

1

2

$$\therefore X = A^{-1} \cdot B = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix} \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

1

$\therefore x = 1, y = 2, z = 3$

1


Using elementary Row operations:

let: $A = IA$

$$\begin{aligned} \left[\begin{array}{ccc} 1 & 2 & 3 \\ 2 & 5 & 7 \\ -2 & -4 & -5 \end{array} \right] &= \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] A && 1 \\ \Rightarrow \left[\begin{array}{ccc} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{array} \right] &= \left[\begin{array}{ccc} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 2 & 0 & 1 \end{array} \right] A &\text{ {Using, } R_2 \rightarrow R_2 - 2R_1; R_3 \rightarrow R_3 + 2R_1} & \left. \right\} \\ \Rightarrow \left[\begin{array}{ccc} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{array} \right] &= \left[\begin{array}{ccc} 5 & -2 & 0 \\ -2 & 1 & 0 \\ 2 & 0 & 1 \end{array} \right] A &\text{ {Using, } R_1 \rightarrow R_1 - 2R_2} & 4 \\ \Rightarrow \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] &= \left[\begin{array}{ccc} 3 & -2 & -1 \\ -4 & 1 & -1 \\ 2 & 0 & 1 \end{array} \right] A &\text{ {Using, } R_1 \rightarrow R_1 - R_3; R_2 \rightarrow R_2 - R_3} & \left. \right\} \\ \therefore A^{-1} &= \left[\begin{array}{ccc} 3 & -2 & -1 \\ -4 & 1 & -1 \\ 2 & 0 & 1 \end{array} \right] && 1 \end{aligned}$$

26.

Correct figure:



Pt. of intersection, $x = 4$

$$\begin{aligned} \text{Area of shaded region} &= \int_0^4 x \, dx + \int_4^{4\sqrt{2}} \sqrt{32 - x^2} \, dx && 1 \\ &= \left. \frac{x^2}{2} \right|_0^4 + \left. \left\{ \frac{x}{2} \sqrt{32 - x^2} + 16 \sin^{-1} \frac{x}{4\sqrt{2}} \right\} \right|_4^{4\sqrt{2}} && 2 \\ &= 8 + 16 \frac{\pi}{2} - 8 - 4\pi = 4\pi && 1 \end{aligned}$$

27. Put $\sin x - \cos x = t$, $(\cos x + \sin x) dx = dt$, $1 - \sin 2x = t^2$

1

$$\left. \begin{array}{l} \text{when } x=0, t=-1 \\ \text{and } x=\pi/4, t=0 \end{array} \right\}$$

 $\frac{1}{2}$

$$\therefore I = \int_0^{\pi/4} \frac{\sin x + \cos x}{16+9\sin 2x} dx = \int_{-1}^0 \frac{1}{16+9(1-t^2)} dt = \int_{-1}^0 \frac{1}{25-9t^2} dt$$

2

$$\Rightarrow I = \frac{1}{30} \log \left| \frac{5+3t}{5-3t} \right|_{-1}^0$$

 $1\frac{1}{2}$

$$= \frac{1}{30} \left[0 - \log \frac{1}{4} \right] = -\frac{1}{30} \log \frac{1}{4} \text{ or } \frac{1}{15} \log 2$$

1

OR

Here $f(x) = x^2 + 3x + e^x$, $a = 1$, $b = 3$, $nh = 2$

1

$$\therefore \int_1^3 (x^2 + 3x + e^x) dx = \lim_{h \rightarrow 0} [f(1) + f(1+h) + \dots + f(1+\overline{n-1}h)]$$

1

$$= \lim_{h \rightarrow 0} \left[4(nh) + \frac{(nh-h)(nh)(2nh-h)}{6} + \frac{5(nh-h)(nh)}{2} + \frac{h}{e^h - 1} \times e \times (e^{nh} - 1) \right]$$

3

$$= 8 + \frac{8}{3} + 10 + e(e^2 - 1) = \frac{62}{3} + e^3 - e$$

1

28. General point on the line is: $(2 + 3\lambda, -1 + 4\lambda, 2 + 2\lambda)$

 $1\frac{1}{2}$

As the point lies on the plane

$$\therefore 2 + 3\lambda + 1 - 4\lambda + 2 + 2\lambda = 5 \Rightarrow \lambda = 0$$

 $1\frac{1}{2}$


$$\therefore \text{Point is } (2, -1, 2)$$

1

$$\text{Distance} = \sqrt{(2 - (-1))^2 + (-1 - (-5))^2 + (2 - (-10))^2} = 13$$

2

29.



Let number of packets of type $A = x$

and number of packets of type $B = y$

\therefore L.P.P. is: Maximize, $Z = 0.7x + y$

1

subject to constraints:

$$\begin{aligned} 4x + 6y &\leq 240 \quad \text{or} \quad 2x + 3y \leq 120 \\ 6x + 3y &\leq 240 \quad \text{or} \quad 2x + y \leq 80 \end{aligned} \quad \left. \right\}$$

2

$$x \geq 0, y \geq 0$$

2

Correct graph

$$Z(0, 0) = 0, Z(0, 40) = 40$$

$$Z(40, 0) = 28, Z(30, 20) = 41 \text{ (Max.)}$$

\therefore Max. profit is ₹ 41 at $x = 30, y = 20$.

1